



HOW TO TEST YOUR DRUGS WITH FENTANYL TEST STRIPS



Prepare your drugs. You only need a small amount. Residue from the bag/cooker works. If you are testing pills, crush and dissolve the whole pill to ensure the test is accurate.

Add about 10 drops of fresh water. The amount of water doesn't need to be exact. It's always better to add more water than less. Mix well.



Dip wavy-line end of test strip in water and allow water to rise up the strip. Be careful to not dip it past the blue line!



Remove and lay on a flat surface. Allow 5 minutes for the results to develop.

TESTING AFTER USE

NOTE:

It's always best to test beforehand, but you can also test the residue on a used cooker – just add water and run the test! Used bags can also be tested by placing them directly into a cooker/ cup

RESULTS

1 LINE = FENTANYL

2 LINES = NO FENTANYL

If the result is not clear, run the test again.



HOW TO TEST YOUR DRUGS WITH FENTANYL TEST STRIPS



Prepare your drugs. You only need a small amount. Residue from the bag/cooker works. If you are testing pills, crush and dissolve the whole pill to ensure the test is accurate.

Add about 10 drops of fresh water. The amount of water doesn't need to be exact. It's always better to add more water than less. Mix well.



Dip wavy-line end of test strip in water and allow water to rise up the strip. Be careful to not dip it past the blue line!



Remove and lay on a flat surface. Allow 5 minutes for the results to develop.

TESTING AFTER USE

NOTE:

It's always best to test beforehand, but you can also test the residue on a used cooker – just add water and run the test! Used bags can also be tested by placing them directly into a cooker/ cup

RESULTS

1 LINE = FENTANYL

2 LINES = NO FENTANYL

If the result is not clear, run the test again.

This process can be used to test powdered drugs, pills, crystals and blotter tabs!

Crystals: Finely grind the sample then run the test as instructed.

PILLS: The entire pill needs to be tested since these mixtures can be uneven. Crush and dissolve the whole pill to protect against the 'chocolate chip cookie effect.'

BLOTTER TABS: Cut off a small piece and then run the test as instructed.

AMPHETAMINE-BASED DRUGS (SPEED, METH, MDMA, ETC): A large amount of these drugs can cause a false positive so only use a few grains of powder and run the test with about a half a cup of water,

This process can be used to test powdered drugs, pills, crystals and blotter tabs!

Crystals: Finely grind the sample then run the test as instructed.

PILLS: The entire pill needs to be tested since these mixtures can be uneven. Crush and dissolve the whole pill to protect against the 'chocolate chip cookie effect.'

BLOTTER TABS: Cut off a small piece and then run the test as instructed.

AMPHETAMINE-BASED DRUGS (SPEED, METH, MDMA, ETC): A large amount of these drugs can cause a false positive so only use a few grains of powder and run the test with about a half a cup of water,

What if I get a positive result?

If you know that fentanyl may be present, you can be prepared with naloxone in case of an overdose.

If you decide to use your drugs after they test positive for fentanyl, here are some tips to stay safe:

- Do a tester shot, go slow
- Have naloxone nearby
- Tell someone you trust so they can check on you & help

A negative result doesn't guarantee safety. No matter the type of result you get, using slowly helps feel out your tolerance and the strength of the batch.

What if I get a positive result?

If you know that fentanyl may be present, you can be prepared with naloxone in case of an overdose.

If you decide to use your drugs after they test positive for fentanyl, here are some tips to stay safe:

- Do a tester shot, go slow
- Have naloxone nearby
- Tell someone you trust so they can check on you & help

A negative result doesn't guarantee safety. No matter the type of result you get, using slowly helps feel out your tolerance and the strength of the batch.

Important things to remember:

- U.S. Drug Enforcement Agency has not found fentanyl in any of its seized cannabis products. There is no evidence that fentanyl is being added to the cannabis supply. Even if fentanyl was added to cannabis, fentanyl burns at a lower temperature than cannabis does, so the flame used to burn the cannabis would likely destroy fentanyl before it could be consumed.
- Know that fentanyl test strips can't 100% guarantee that fentanyl is not present. They can't test for fentanyl analogues and can't tell you how much fentanyl is in your drugs.
- Fentanyl isn't the only thing you need to look out for. Drugs are cut with many things like other drugs, Tylenol, or tranquilizers. Use caution often.

Important things to remember:

- U.S. Drug Enforcement Agency has not found fentanyl in any of its seized cannabis products. There is no evidence that fentanyl is being added to the cannabis supply. Even if fentanyl was added to cannabis, fentanyl burns at a lower temperature than cannabis does, so the flame used to burn the cannabis would likely destroy fentanyl before it could be consumed.
- Know that fentanyl test strips can't 100% guarantee that fentanyl is not present. They can't test for fentanyl analogues and can't tell you how much fentanyl is in your drugs.
- Fentanyl isn't the only thing you need to look out for. Drugs are cut with many things like other drugs, Tylenol, or tranquilizers. Use caution often.

Sometimes opioid overdoses can still happen.

Have naloxone? Use it!

- Administer & remember to provide rescue breathing if they aren't breathing.

Don't have naloxone?

- Rescue breathing: Breathe one breath every 3 seconds until they awake & breathe without help. Lack of oxygen can cause severe brain damage if not breathing for too long.
- Sternum Rub: If someone is unconscious but still breathing, use your knuckles to rub the middle of their chest (sternum).

Sometimes opioid overdoses can still happen.

Have naloxone? Use it!

- Administer & remember to provide rescue breathing if they aren't breathing.

Don't have naloxone?

- Rescue breathing: Breathe one breath every 3 seconds until they awake & breathe without help. Lack of oxygen can cause severe brain damage if not breathing for too long.
- Sternum Rub: If someone is unconscious but still breathing, use your knuckles to rub the middle of their chest (sternum).